

# PP2A methylation controls sensitivity and resistance to $\beta$ -amyloid-induced cognitive and electrophysiological impairments

Russell E. Nicholls<sup>a,b,1</sup>, Jean-Marie Sontag<sup>c</sup>, Hong Zhang<sup>a,b</sup>, Agnieszka Staniszewski<sup>a,b</sup>, Shijun Yan<sup>a,b</sup>, Carla Y. Kim<sup>d</sup>, Michael Yim<sup>d</sup>, Caitlin M. Woodruff<sup>d</sup>, Erland Arning<sup>e</sup>, Brandi Wasek<sup>e</sup>, Deqi Yin<sup>d,f</sup>, Teodoro Bottiglieri<sup>g</sup>, Estelle Sontag<sup>c</sup>, Eric R. Kandel<sup>d,f,g,1</sup>, and Ottavio Arancio<sup>a,b</sup>

<sup>a</sup>Department of Pathology and Cell Biology, Columbia University, New York, NY 10032; <sup>b</sup>The Taub Institute for Research on Alzheimer's Disease and the Aging Brain, Columbia University, New York, NY 10032; <sup>c</sup>The School of Biomedical Sciences and Pharmacy, Faculty of Health and Medicine, Hunter Medical Research Institute, The University of Newcastle, Callaghan, NSW 2308, Australia; <sup>d</sup>Department of Neuroscience, Columbia University, New York, NY 10032; <sup>e</sup>Institute of Metabolic Disease, Baylor Research Institute, Dallas, TX 75226; <sup>f</sup>Howard Hughes Medical Institute, Columbia University, New York, NY 10032; and <sup>g</sup>Kavli Institute for Brain Science, Columbia University, New York, NY 10032

Contributed by Eric R. Kandel, February 4, 2016 (sent for review October 23, 2015; reviewed by Domenico Pratico and David M. Virshup)

Elevated levels of the  $\beta$ -amyloid peptide (A $\beta$ ) are thought to contribute to cognitive and behavioral impairments observed in Alzheimer's disease (AD). Protein phosphatase 2A (PP2A) participates in multiple molecular pathways implicated in AD, and its expression and activity are reduced in postmortem brains of AD patients. PP2A is regulated by protein methylation, and impaired PP2A methylation is thought to contribute to increased AD risk in hyperhomocysteinemic individuals. To examine further the link between PP2A and AD, we generated transgenic mice that overexpress the PP2A methyltransferase, protein phosphatase methyltransferase-1 (PME-1), or the PP2A methyltransferase, leucine carboxyl methyltransferase-1 (LCMT-1), and examined the sensitivity of these animals to behavioral and electrophysiological impairments caused by exogenous A $\beta$  exposure. We found that PME-1 overexpression enhanced these impairments, whereas LCMT-1 overexpression protected against A $\beta$ -induced impairments. Neither transgene affected A $\beta$  production or the electrophysiological response to low concentrations of A $\beta$ , suggesting that these manipulations selectively affect the pathological response to elevated A $\beta$  levels. Together these data identify a molecular mechanism linking PP2A to the development of AD-related cognitive impairments that might be therapeutically exploited to target selectively the pathological effects caused by elevated A $\beta$  levels in AD patients.

Alzheimer's disease | protein phosphatase 2A | methylation |  $\beta$ -amyloid | cognitive impairment

Multiple observations suggest a role for the serine/threonine protein phosphatase 2A (PP2A) in the molecular pathways that underlie Alzheimer's disease (AD). Analyses conducted on postmortem AD brains have found reduced PP2A expression and activity, and studies conducted in animal models have found that inhibiting PP2A produces AD-like tau pathology and cognitive impairment (1–3). One of the ways in which PP2A may affect AD is through its role as the principal tau phosphatase (4–7). PP2A also interacts with a number of kinases implicated in AD including glycogen synthase kinase 3 $\beta$  (GSK3 $\beta$ ), cyclin-dependent kinase 5 (CDK5), and ERK and JNK as well as amyloid precursor protein and the NMDA and metabotropic glutamate receptors (reviewed in ref. 2).

PP2A is a heterotrimeric protein composed of a catalytic, scaffolding, and regulatory subunit. Each subunit is encoded by multiple genes and splice isoforms, and the subunit composition of a particular PP2A molecule determines its subcellular distribution and substrate specificity (reviewed in ref. 2). One of the ways in which PP2A activity is regulated is through C-terminal methylation of the catalytic subunit (reviewed in refs. 8 and 9). Impaired methyl-donor metabolism is a risk factor for AD (10, 11), and PP2A dysregulation caused by impaired methylation is thought to be one of the molecular mechanisms contributing to this increased risk (12–14). Methylation promotes the formation of PP2A holoenzymes that

contain B $\alpha$  regulatory subunits (7, 13, 15–19), and these forms of PP2A exhibit the greatest tau phosphatase activity (6, 7).

PP2A methylation is catalyzed *in vivo* by the methyl transferase, leucine carboxyl methyltransferase 1 (LCMT-1) (20–22), and its demethylation is catalyzed by the methyltransferase, protein phosphatase methyltransferase 1 (PME-1) (23–25). To explore the role of PP2A in AD further, we generated lines of transgenic mice that overexpress these enzymes and tested their effect on the sensitivity of animals to electrophysiological and behavioral impairments caused by  $\beta$ -amyloid (A $\beta$ ). We found that LCMT-1 overexpression protected animals from A $\beta$ -induced impairments, whereas overexpression of PME-1 worsened A $\beta$  neurotoxicity. Neither transgene affected endogenous A $\beta$  levels, suggesting that they acted by altering the response to A $\beta$  rather than A $\beta$  production. We also found that PME-1 and LCMT-1 overexpression were without effect on the electrophysiological response to picomolar A $\beta$  application, suggesting that they selectively affected the response to pathological A $\beta$  concentrations. Together these data indicate that this pathway has potential as a therapeutic avenue for AD that acts not by targeting A $\beta$  production but by selectively altering the response to pathological levels of A $\beta$ .

## Results

**PME-1 Overexpression in Transgenic Mice.** To test the effect of reduced PP2A methylation on the sensitivity to A $\beta$ -induced

### Significance

Elevated levels of the  $\beta$ -amyloid peptide (A $\beta$ ) are thought to contribute to the cognitive impairments associated with Alzheimer's disease (AD). We found that by genetically targeting the methylation of protein phosphatase 2A (PP2A) in transgenic mice, we could alter the sensitivity of animals to electrophysiological and cognitive impairments caused by A $\beta$  exposure without affecting A $\beta$  production or the electrophysiological response to low concentrations of A $\beta$ . These data support a role for PP2A methylation in contributing to AD risk and identify a potential therapeutic pathway that might be exploited to target the pathological actions of A $\beta$  selectively.

Author contributions: R.E.N., J.-M.S., E.S., E.R.K., and O.A. designed research; R.E.N., J.-M.S., H.Z., A.S., S.Y., C.Y.K., M.Y., C.M.W., E.A., B.W., and D.Y. performed research; R.E.N., T.B., and E.R.K. contributed new reagents/analytic tools; R.E.N., J.-M.S., H.Z., A.S., S.Y., T.B., E.S., and O.A. analyzed data; and R.E.N. and E.S. wrote the paper.

Reviewers: D.P., Temple University; and D.M.V., Duke-National University of Singapore Graduate Medical School.

The authors declare no conflict of interest.

<sup>1</sup>To whom correspondence may be addressed. Email: erk5@columbia.edu or rn95@columbia.edu.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1521018113/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1521018113/-DCSupplemental).


impairments, we generated mice carrying a transgene encoding FLAG epitope-tagged murine PP2A methyltransferase (PME-1) under the control of a synthetic *tetO* promoter. To drive expression of the *PME-1* transgene in neurons in the forebrain, we crossed these animals with mice carrying a second transgene that was under the control of a promoter region from the calcium calmodulin kinase II $\alpha$  (*CaMKII $\alpha$* ) gene and expressed a synthetic tetracycline-responsive transactivator (tTA) (26). In this system, *PME-1* transgene expression is activated when tTA binds to the *tetO* promoter in cells where tTA expression is driven by the *CaMKII $\alpha$*  promoter (Fig. 1A). Although not used in the current experiments, this system also affords the ability to suppress *tetO* promoter-driven transgene expression through doxycycline administration. *PME-1* transgene-specific RNA in situ hybridization revealed expression throughout the forebrain, including the striatum, olfactory bulb, cortex, and hippocampus, in animals that carried both the *tetO-PME* and *CaMK-tTA* transgenes but not in single-transgenic siblings (Fig. 1B, Left). Immunohistochemistry on brain sections from these animals using an antibody specific to the FLAG epitope tag detected transgenic protein in cell bodies and dendrites of pyramidal cells of the hippocampal CA1 region of double-transgenic animals that was not present in single-transgenic control animals (Fig. 1B, Right).

Quantitative Western blot analysis of hippocampal extracts from *tetO-PME/CaMKtTA* double-transgenic animals revealed a significant increase in PME-1 expression compared with single-transgenic control animals (Fig. 1C). Transgene expression in these animals did not affect PP2A catalytic subunit expression, PP2A B $\alpha$  regulatory subunit expression, LCMT-1 expression (Fig. 1C), or levels of the methylation metabolites S-adenosylmethionine (SAM) and S-adenosylhomocysteine (SAH) (Fig. S1). Quantitative Western blot analysis did reveal a significant reduction in methylated PP2A levels in animals overexpressing PME-1 as compared with controls (Fig. 1D). This change was accompanied by an increase in tau phosphorylation at sites reported to be targets of PP2A (5, 13), but not in total tau levels (Fig. 1E), as well as increased phosphorylation of amyloid precursor protein (APP) at threonine 668 (Fig. 1F), which also is a target of PP2A (13). ELISA measurements of A $\beta$ <sub>1-40</sub> and A $\beta$ <sub>1-42</sub> levels in hippocampal homogenates from animals overexpressing PME-1 and control animals revealed no significant change in basal levels of either peptide (Fig. 1H).

**PME-1 Overexpression Increases Sensitivity to Behavioral and Physiological Impairments Caused by Exposure to Subthreshold Doses of Oligomeric A $\beta$  but Not to Picomolar A $\beta$  Doses.** Mice overexpressing PME-1 were overtly indistinguishable from their single-transgenic control siblings. They were fertile and were recovered at the expected frequencies from crosses of double-transgenic males to wild-type females (Fig. S2). Analysis of these animals' behavior in a novel open field revealed no genotype effects (Fig. S3).


To test the effect of PME-1 overexpression on the behavioral impairments that result from acute A $\beta$  exposure, we tested these animals in a hippocampus-dependent contextual fear-conditioning task previously shown to be sensitive to A $\beta$  administration (27–29). Vehicle-treated animals overexpressing PME-1 exhibited a level of freezing 24 h after training in this task similar to that of vehicle-treated single-transgenic control siblings (Fig. 2A). Control animals infused with a subthreshold dose of oligomeric A $\beta$  (bilateral infusions of 1  $\mu$ L of 75 nM oligomeric A $\beta$ ) also exhibited levels of freezing similar to those in the vehicle-treated controls. However, animals overexpressing PME-1 that received a subthreshold dose of A $\beta$  exhibited significantly less freezing than the other three groups, suggesting that animals that overexpress PME-1 are more sensitive to A $\beta$ -induced cognitive impairment in this task. We observed no differences in baseline freezing, open-field ambulatory activity, or shock perception among these groups, suggesting that these differences in contextual fear conditioning did not result from differences in baseline behaviors or sensory perception (Fig. 2A and Fig. S4).

As a second test of the effect of PME-1 overexpression on A $\beta$ -induced cognitive impairments, we tested these animals on a 2-d radial arm water maze task (29, 30) and found that



**Fig. 1.** *PME-1* transgene expression in the forebrain of transgenic mice. (A) Diagram of the *tetO/tTA* system in which FLAG-tagged *PME-1* transgene expression is driven by the tTA transactivator under the control of a *CaMKII $\alpha$*  promoter fragment. (B) Representative images of transgene-specific RNA in situ hybridization of whole-brain sagittal sections (Left) and immunofluorescent images of anti-FLAG-tag staining in hippocampal CA1 region pyramidal cells (Right) from animals with the indicated genotypes. (C) Western blot analyses of *PME-1*, PP2A catalytic subunit (PP2A/C), and regulatory (B $\alpha$ ) subunit and LCMT-1 protein expression in hippocampal homogenates carried out using primary antibodies directed against the proteins indicated at left and normalized to  $\beta$ -actin reveal a 379  $\pm$  44% increase in PME expression in *tetO-PME/CaMKtTA* animals (upper transgenic + lower endogenous bands) vs. *tetO-PME* or *CaMKtTA* controls ( $P < 0.001$ ) and unchanged levels of PP2A/C and B $\alpha$  (103  $\pm$  1.4 and 98  $\pm$  2.7% of control, respectively;  $P > 0.05$ ). (D) Western blot analyses of PP2A methylation in hippocampal homogenates from transgenic animals overexpressing *PME-1* carried out using a methyl-PP2A/C-specific and a methylation-insensitive total PP2A/C antibody reveal a 54.8  $\pm$  6.6% reduction in methylated PP2A levels in *tetO-PME/CaMKtTA* animals vs. controls;  $P < 0.001$ . (E) Western blots showing increased tau phosphorylation but normal tau expression in hippocampal homogenates from transgenic animals overexpressing *PME-1* were conducted using the phospho-specific tau antibodies PHF-1, S262, and AT8 or total tau and normalized to total tau or  $\beta$ -actin as indicated (PHF-1: Ser396/Ser-404: 210  $\pm$  7%,  $P < 0.001$ ; AT8: Ser202/Thr205: 139  $\pm$  15%,  $P < 0.05$ ; anti-phospho-S262: 116  $\pm$  7.5%,  $P < 0.05$ ; total tau: 98  $\pm$  5.4%,  $P > 0.05$  compared with controls). (F) Western blots using phospho-Thr-668-specific and total anti-APP antibodies reveal a 150  $\pm$  5% increase in APP phosphorylation in hippocampal homogenates from transgenic animals overexpressing *PME-1* compared with controls;  $P < 0.001$ . (G) Graph of the mean percent of control values ( $\pm$  SEM) obtained for the blots shown in C–F. (H) Graphs of the values and means ( $\pm$  SEM) obtained by ELISA for A $\beta$ <sub>1-40</sub> (Left) or A $\beta$ <sub>1-42</sub> (Right) conducted on hippocampal homogenates from animals overexpressing *PME-1* ( $n = 9$ ) and controls ( $n = 10$ ) and normalized to the total amount of protein in each sample;  $P > 0.05$  for A $\beta$ <sub>1-40</sub> and A $\beta$ <sub>1-42</sub> *PME* vs. control comparisons.

*PME-1* overexpression also sensitized animals to A $\beta$ -induced impairments in this task. *PME-1* overexpression did not affect the performance of vehicle-infused animals, and a subthreshold dose of A $\beta$  did not affect the performance of control animals. However, a subthreshold dose of A $\beta$  did significantly impair the performance of animals overexpressing *PME-1* (Fig. 2B). Tests



**Fig. 2.** PME-1 overexpression increases behavioral and electrophysiological impairments caused by subthreshold doses of A $\beta$  without affecting picomolar A $\beta$  responses. (A) Average percent of time ( $\pm$  SEM) spent freezing during the initial exposure to the training context (baseline) and 24 h after foot shock for the indicated genotype and treatment groups ( $n = 13$  animals per group). Two-way ANOVA effect of training:  $F(1,48) = 138.45$ ,  $P < 0.0001$ ; training  $\times$  group interaction:  $F(3,48) = 3.43$ ,  $P < 0.05$ . Bonferroni post hoc comparisons of freezing at 24 h for treated animals overexpressing PME-1 vs. all other groups were  $P < 0.01$  and  $P > 0.05$  for baseline freezing responses. (B) Average number of errors committed ( $\pm$  SEM) during each three-trial training block of a 2-d radial arm water maze task for the indicated genotype and treatment groups. Two-way repeated-measures ANOVA with block and treatment as factors for PME vehicle vs. PME + A $\beta$  on day 2 (blocks 6–10):  $F(1,24) = 20.81$ ,  $P < 0.0001$  for treatment;  $P > 0.05$  for control vehicle vs. PME vehicle and for control vehicle vs. control A $\beta$  on day 2.  $n = 13$  animals per group. (C and D) Time course of averaged Schaffer collateral fEPSP responses in slices treated with vehicle (C) or 50 nM A $\beta$  (D) 20 min before delivery of theta-burst stimulation (arrow). Repeated-measures ANOVA for PME + A $\beta$  vs. control + A $\beta$ :  $F(1,22) = 5.23$ ,  $P < 0.05$ ;  $P > 0.05$  for control vehicle vs. PME vehicle and for control vehicle vs. control A $\beta$ .  $n > 10$  slices per group. (E) Dose–response curve showing increased A $\beta$ -induced LTP inhibition in animals overexpressing PME-1 compared with controls. Plotted are average potentiated responses ( $\pm$  SEM) measured 105–115 min after theta-burst stimulation in slices treated with oligomeric A $\beta$  at the indicated concentration 20 min before potentiating stimulation. Two-way ANOVA for genotype:  $F(1,132) = 5.42$ ,  $P < 0.05$ . (F) Time course of averaged Schaffer collateral fEPSP responses ( $\pm$  SEM) in hippocampal slices from animals overexpressing PME-1 and control animals treated with vehicle or with 200 pM A $\beta$  20 min before delivery of theta-burst stimulation (arrow). Repeated measures ANOVA: PME + A $\beta$  vs. PME + vehicle:  $F(1,15) = 5.893$ ,  $P < 0.05$ ; PME + A $\beta$  vs. control + A $\beta$ :  $F(1,20) = 0.6677$ ,  $P > 0.05$ .  $n = 8$  PME + vehicle, 9 PME + A $\beta$ , 11 control + vehicle, and 13 control + A $\beta$  slices.

of these animals on a visible platform version of the Morris water maze revealed no differences in escape latency or swim speed among these groups, suggesting that the combination of PME-1 overexpression and subthreshold A $\beta$  exposure does not measurably impact motor performance, perception, or motivation in a nonspatial water maze task (Fig. S4).


Activity-dependent changes in the efficacy of synaptic transmission within the hippocampus are thought to be required for particular forms of memory, and interference with these changes, caused by elevated levels of A $\beta$ , is thought to contribute to

AD-associated memory impairments (31). Because PME-1 overexpression enhances A $\beta$ -induced memory impairments, we sought to determine whether PME-1 overexpression also might enhance A $\beta$ -induced impairment of activity-dependent synaptic plasticity. To do so, we tested the ability of A $\beta$  to impair theta-burst-induced long-term potentiation (LTP) at Schaffer collateral synapses (32–34). This protocol produced robust potentiation that was comparable in animals overexpressing PME-1 and controls, suggesting that PME-1 overexpression alone did not affect LTP under these conditions (Fig. 2C). However, bath application of a subthreshold dose of A $\beta$  (50 nM), which produced no significant effect on LTP in slices from control animals, significantly impaired LTP in slices from animals overexpressing PME-1 (Fig. 2C and D). This increase in the sensitivity of slices overexpressing PME-1 to A $\beta$ -induced LTP impairment also could be seen as a shift in the relationship between A $\beta$  concentration and the corresponding LTP impairment (Fig. 2E). This enhancement of A $\beta$ -induced LTP impairment was not accompanied by a shift in the stimulus intensity/response relationship in these animals (Fig. S4) and parallels the behavioral data we describe above, suggesting that PME-1-mediated enhancement of A $\beta$ 's effects on synaptic plasticity may lead to the enhanced A $\beta$ -induced behavioral impairments we observed.

The impairment of LTP by A $\beta$  is thought to reflect a pathological process that occurs at high A $\beta$  concentrations in patients with AD. However, at low (picomolar) concentrations of A $\beta$ , LTP is enhanced, and this effect is thought to reflect a normal physiological function of the protein (33, 35, 36). To determine whether PME overexpression also affects sensitivity to picomolar concentrations of A $\beta$ , we recorded theta-burst-evoked LTP at Schaffer collateral synapses in the presence or absence of 200 pM oligomeric A $\beta$ . In these experiments, picomolar A $\beta$  application caused a significant enhancement of LTP over corresponding vehicle-treated slices that was comparable in both control animals and animals overexpressing PME-1 (Fig. 2F). These results are consistent with the observation that baseline LTP, behavior, and A $\beta$  production are all normal in mice overexpressing PME-1 and suggest that PME-1 overexpression may selectively affect the response of these animals to pathological levels of A $\beta$  without affecting A $\beta$ 's normal physiological function.

**LCMT-1 Overexpression in Transgenic Mice.** Because we found that PME-1 overexpression sensitized animals to A $\beta$ -induced impairments, we sought to determine whether LCMT-1 overexpression might exert an opposite effect on A $\beta$  sensitivity. To do so, we generated mice carrying a transgene encoding FLAG epitope-tagged murine PP2A methyltransferase (LCMT-1) under the control of a synthetic *tetO* promoter and crossed these animals with mice carrying the same *CaMK- $\alpha$*  transgene. *LCMT-1* transgene-specific RNA in situ hybridization revealed expression throughout the forebrain, including the striatum, olfactory bulb, cortex, and hippocampus, in animals that carried both the *tetO-LCMT* and *CaMK- $\alpha$*  transgenes but not in single-transgenic siblings (Fig. 3A, Left). Anti-FLAG epitope tag immunohistochemistry on brain sections from *tetO-LCMT/CaMK $\alpha$*  double-transgenic animals revealed transgenic protein in cell bodies and dendrites of pyramidal cells of the hippocampal CA1 region of double-transgenic animals that was not present in single-transgenic control animals (Fig. 3A, Right).

Quantitative Western blot analysis of hippocampal extracts from *tetO-LCMT/CaMK $\alpha$*  double-transgenic animals revealed a significant increase in *LCMT-1* expression compared with single-transgenic control animals (Fig. 3B). As was the case for PME-1 overexpression, *LCMT-1* transgene expression in these animals did not affect PP2A catalytic subunit expression, PP2A B $\alpha$  regulatory subunit expression (Fig. 3B), or methylation metabolite levels (Fig. S1), nor did it affect endogenous PME-1 expression (Fig. 3B). Western blot analysis using methyl-PP2A/C-specific antibodies failed to reveal any significant change in PP2A methylation in *tetO-LCMT/CaMK $\alpha$*  double-transgenic animals (Fig. 3C). However, this result likely reflects the nearly saturating levels of PP2A methylation that exist under basal conditions (37). In addition, with the exception of a modest decrease in paired helical



**Fig. 3.** *LCMT-1* transgene expression in the forebrain of transgenic mice. (A) Representative images of transgene-specific RNA in situ hybridization of whole-brain sagittal sections (*Left*) and immunofluorescent images of anti-FLAG-tag staining in hippocampal CA1 region pyramidal cells (*Right*) from animals with the indicated genotypes. (B) Western blot analyses of *LCMT-1*, *PP2A/C*, regulatory ( $B\alpha$ ) subunit, and *PME-1* protein expression in hippocampal homogenates from *tetO-LCMT/CaMKtTA* double-transgenic animals and single-transgenic siblings carried out using primary antibodies directed against the proteins indicated at left and normalized to  $\beta$ -actin reveal a  $152 \pm 13\%$  increase in *LCMT* expression in *tetO-LCMT/CaMKtTA* animals (upper transgenic + lower endogenous bands) vs. *tetO-LCMT* or *CaMKtTA* controls ( $P < 0.001$ ) and unchanged levels of *PP2A/C* and  $B\alpha$  ( $104 \pm 1.0$  and  $104 \pm 2.0\%$  of control, respectively;  $P > 0.05$ ). (C) Western blot analyses of *PP2A* methylation in hippocampal homogenates from transgenic animals overexpressing *LCMT-1* carried out using a methyl-*PP2A/C*-specific and a methylation-insensitive total *PP2A/C* antibody reveal no change ( $105.9 \pm 7.3\%$  of control) in methylated *PP2A* levels in *tetO-LCMT/CaMKtTA* animals vs. controls;  $P > 0.05$ . (D) Western blots showing tau phosphorylation and total tau expression in hippocampal homogenates from transgenic animals overexpressing *LCMT-1* conducted using phospho-specific tau antibodies PHF-1, S262, and AT8 or total tau and normalized to total tau or  $\beta$ -actin as indicated. PHF-1:  $79.5 \pm 12.4\%$  of control,  $P < 0.05$ ; AT8:  $95 \pm 2.9\%$  of control,  $P > 0.05$ ; S262:  $95 \pm 2.4\%$  of control,  $P > 0.05$ ; total tau:  $106 \pm 6.7\%$  of control,  $P > 0.05$ . (E) Western blots showing reduced APP phosphorylation ( $56 \pm 5\%$  of control) in hippocampal homogenates from transgenic animals overexpressing *LCMT-1* conducted using a phospho-Thr-668 antibody (*Upper*) and a phosphorylation-insensitive total APP antibody for normalization (*Lower*);  $P < 0.001$ . (F) Graph of the mean percent of control values ( $\pm$  SEM) obtained for the blots shown in *B–E*. (G) Graphs of the values and means ( $\pm$  SEM) obtained by ELISA for  $A\beta_{1-40}$  (*Left*) or  $A\beta_{1-42}$  (*Right*) conducted on hippocampal homogenates from animals overexpressing *LCMT* ( $n = 8$ ) and controls ( $n = 13$ ) and normalized to the total amount of protein in each sample.  $P > 0.05$  for  $A\beta_{1-40}$  and  $A\beta_{1-42}$  *LCMT* vs. control comparisons.

filament 1 (PHF-1) antibody immunoreactivity, we did not observe corresponding decreases in phospho-tau immunoreactivity in animals overexpressing *LCMT-1* (Fig. 3D). However, this result also may be caused by the relatively low levels of phosphorylation that exist at these sites under basal conditions (38). We did find that *LCMT-1* overexpression led to a decrease in APP phosphorylation at the *PP2A*-sensitive Thr-668 site (Fig. 3E). Like *PME-1* overexpression, *LCMT-1* overexpression was without significant effect on basal hippocampal  $A\beta_{1-40}$  or  $A\beta_{1-42}$  levels (Fig. 3F).

### **LCMT-1 Overexpression Reduces Physiological and Behavioral Impairments Caused by Exposure to Nanomolar Concentrations of $A\beta$ Without Affecting Responses to Picomolar Concentrations of $A\beta$ .**


Like the mice overexpressing *PME-1*, animals overexpressing *LCMT-1* were overtly indistinguishable from their single-transgenic control siblings. They were fertile and were recovered at the expected frequencies from crosses of double-transgenic males to wild-type females (Fig. S2). Analysis of these animals' behavior in a novel open field revealed no genotype effects (Fig. S5). To determine whether *LCMT-1* overexpression might protect against  $A\beta$ -induced impairments, we subjected animals overexpressing *LCMT-1* to the contextual fear-conditioning and radial arm water maze tasks described above. In these experiments, however, we infused animals with  $A\beta$  at a higher concentration that produced behavioral impairments in control animals (bilateral infusions of  $1 \mu\text{L}$  of  $200 \text{ nM}$  oligomeric  $A\beta$ ).

We found that *LCMT-1* overexpression protected against  $A\beta$ -induced impairment in the contextual fear-conditioning task. Vehicle-treated control animals exhibited a robust freezing response when tested 24 h after training that was significantly reduced in  $A\beta$ -infused controls but was unaffected in similarly infused animals overexpressing *LCMT-1* (Fig. 4A). This effect was specific to the  $A\beta$ -induced impairment in these animals, because vehicle-treated animals overexpressing *LCMT-1* showed levels of freezing comparable to those of vehicle-treated controls. Moreover, we observed no differences in baseline freezing, open-field ambulatory activity, or shock perception among these groups (Fig. 4A and Fig. S6), suggesting that contextual fear-conditioning performance was not affected by differences in baseline behaviors or sensory perception.

*LCMT-1* overexpression also protected against  $A\beta$ -induced behavioral impairment in the radial arm water maze task.  $A\beta$  significantly impaired performance in control animals but not in similarly infused animals overexpressing *LCMT-1* (Fig. 4B). The performance of vehicle-treated animals overexpressing *LCMT-1* was comparable to that of vehicle-treated controls, suggesting that this difference was specific to the  $A\beta$  response of these animals. Tests on a visible platform version of the Morris water maze revealed no differences in escape latency or swim speed among these groups, suggesting that neither transgene expression nor  $A\beta$  treatment measurably affected motor performance, perception, or motivation in a nonspatial water maze task (Fig. S6).

To test whether *LCMT-1* overexpression also protected against  $A\beta$ -induced LTP impairment, we recorded theta-burst-induced LTP in  $A\beta$ -treated or vehicle-treated acute hippocampal slice preparations from these animals and controls. Bath application of  $A\beta$  at a dose that significantly impaired LTP in single-transgenic controls produced significantly less impairment in animals overexpressing *LCMT-1* (Fig. 4C and D). This effect was not accompanied by changes in the stimulus/response relationship in these animals (Fig. S6). As was the case for mice overexpressing *PME-1*, this correlation between *LCMT*'s behavioral and electrophysiological effects suggests that *LCMT-1* overexpression may protect against  $A\beta$ -induced behavioral impairments by reducing  $A\beta$ -induced electrophysiological impairments.

Because *LCMT-1* overexpression reduced the sensitivity of animals to impairments caused by nanomolar concentrations of  $A\beta$ , we sought to determine whether it also reduced the response to low, picomolar concentrations of  $A\beta$ . To do this, we recorded theta-burst-evoked LTP at Schaffer collateral synapses in the presence or absence of  $200 \text{ pM}$  oligomeric  $A\beta$  and found that LTP enhancement was comparable in control animals and animals overexpressing *LCMT-1* (Fig. 4E). As was the case for animals overexpressing *PME-1*, these results are consistent with the observation that baseline LTP, behavior, and  $A\beta$  production are all normal in mice overexpressing *LCMT-1* and indicate that *LCMT-1* overexpression may protect selectively against the pathological actions of  $A\beta$  without affecting  $A\beta$ 's normal physiological functions.



**Fig. 4.** LCMT-1 overexpression reduces behavioral and electrophysiological impairments caused by oligomeric A $\beta$ . (A) Average percent of time spent freezing ( $\pm$  SEM) during initial exposure to the training context (baseline) and 24 h after foot shock for the indicated genotype and treatment groups. Two-way ANOVA effect of training:  $F(1,43) = 134.51$ ,  $P < 0.0001$ ; training  $\times$  group interaction:  $F(3,43) = 5.50$ ,  $P < 0.01$ . Bonferroni post hoc comparisons of freezing at 24 h:  $P < 0.01$  for A $\beta$ -treated vs. vehicle-treated controls and  $P > 0.05$  for LCMT + A $\beta$  vs. vehicle-treated controls.  $P > 0.05$  for all baseline freezing response comparisons ( $n = 11$  or 12 animals per group). (B) Average number of errors committed ( $\pm$  SEM) during each three-trial training block of a 2-d radial arm water maze task for the indicated genotype and treatment groups ( $n = 11$  or 12 animals per group). Two-way repeated-measures ANOVA with block and treatment as factors for control + vehicle vs. control + A $\beta$  on day 2 (blocks 6–10):  $F(1,21) = 27.74$ ;  $P < 0.0001$  for treatment;  $P > 0.05$  for control + vehicle vs. LCMT + A $\beta$  and control vehicle vs. LCMT vehicle. (C and D) Time course of averaged Schaffer collateral fEPSP responses in slices treated with vehicle (C;  $n = 9$  LCMT, 11 control slices) or 100 nM A $\beta$  (D;  $n = 9$  LCMT, 13 control slices) 20 min before delivery of theta-burst stimulation (arrow). Repeated-measures ANOVA for treatment on vehicle vs. A $\beta$ -treated controls:  $F(1,22) = 10.73$ ;  $P < 0.01$ . Repeated-measures ANOVA for genotype on A $\beta$ -treated LCMT vs. control slices:  $F(1,20) = 5.13$ ,  $P < 0.05$ . (E) Time course of averaged Schaffer collateral fEPSP responses ( $\pm$  SEM) in hippocampal slices from animals overexpressing LCMT-1 and control animals treated with vehicle or 200 pM A $\beta$  20 min before the delivery of theta-burst stimulation (arrow) ( $n = 12$  LCMT + vehicle, 15 LCMT + A $\beta$  slices). Repeated-measures ANOVA for treatment: LCMT + A $\beta$  vs. vehicle:  $F(1,25) = 8.125$ ,  $P = 0.0086$ ; Repeated-measures ANOVA for genotype: LCMT + A $\beta$  vs. control + A $\beta$   $F(1,26) = 0.4638$ ,  $P = 0.5019$ . Pooled control data are plotted in both Fig. 2E and here for comparison.

## Discussion

Here we show that PME-1 overexpression sensitizes mice to electrophysiological and behavioral impairments caused by acute exposure to nanomolar concentrations of synthetic, oligomeric A $\beta$  and that LCMT-1 overexpression protects against these impairments. These complementary effects of PME-1 and LCMT-1 overexpression on A $\beta$  sensitivity are consistent with the opposing actions of these enzymes on PP2A methylation (8, 9) and support the contention that altered PP2A methylation underlies this altered A $\beta$  sensitivity. However, we cannot rule out the possibility that PME-1 and LCMT-1 act on PP2A via mechanisms that do not require methyltransferase or methyltransferase activity, such as PME-mediated displacement of manganese ions from the PP2A catalytic subunit (25) or stabilization of inactive forms of PP2A (39, 40). LCMT-1 and PME-1 also may interact with as yet unidentified substrates.

Although PP2A is the only known substrate for these enzymes, a recent study did find that the closely related phosphatase, PP4, can interact with a mutant form of PME-1 when overexpressed in cultured cells (41).

The effects of PME-1 and LCMT-1 overexpression on A $\beta$  sensitivity exhibit two features with particular relevance for the potential of this pathway as a therapeutic target for AD treatment or prevention. First, the overexpression of PME-1 and LCMT-1 alters the response of cells and animals to A $\beta$ . Although this altered A $\beta$  sensitivity might have been explained by altered levels of endogenous A $\beta$  that shift the threshold for exogenous A $\beta$ -induced impairment, we detected no changes in basal A $\beta_{40}$  and A $\beta_{42}$  levels in brain homogenates from these animals, no changes in baseline LTP or behavioral performance, and no evidence for occlusion of the electrophysiological response of these animals to picomolar A $\beta$  exposure. This effect on A $\beta$  sensitivity is a clear distinction from therapeutic strategies that seek to alter A $\beta$  production or clearance. The second notable feature of this pathway is that it appears to alter selectively the response to pathological concentrations of A $\beta$ . PME-1 and LCMT-1 bidirectionally altered both the electrophysiological and behavioral responses to nanomolar A $\beta$  administration, without affecting baseline LTP, behavioral performance, or the ability of the administration of 200 pM A $\beta$  to enhance LTP. Given that normal physiological levels of A $\beta$  play an important role in regulating neuronal activity (35, 42–44), this ability to target A $\beta$ 's pathological actions selectively could provide an approach that reduces AD-related impairments without interfering with normal nervous system function. Identifying the molecules acting downstream of PP2A in this process also will help clarify the molecular mechanisms by which A $\beta$  exerts its pathological effects. The transgene-dependent changes in tau and APP phosphorylation that we describe mark these molecules as candidates in this regard; however, additional work is needed to determine the relevance of these changes for A $\beta$  sensitivity.

Together these data add to the growing body of evidence suggesting that PP2A-centric therapies may be beneficial for AD and other tauopathies (reviewed in refs. 3 and 45) and specifically highlight the PP2A-regulatory enzymes PME-1 and LCMT-1 as potential therapeutic targets for this purpose.

## Methods

**Transgenic Animals.** *tetO* promoter-driven constructs expressing either Flag-tagged murine PME-1 or Flag-tagged murine LCMT-1 were generated using standard molecular cloning techniques and were used to generate transgenic mice by pronuclear injection into C57BL/6J oocytes. Transgene-containing animals were crossed to an existing *CaMKII $\alpha$ -tTA* line (26), also in a C57BL/6J background, and double-transgenic animals were outcrossed to wild-type 129SVEV/TAC mice to generate the C57BL/6J  $\times$  129SVEV/TAC F1 animals used for experiments. All experiments were carried out in a manner consistent with National Institutes of Health guidelines (46) and were approved by the Columbia University Institutional Animal Care and Use Committee. We used adult animals between 3 and 6 mo of age for all experiments. Males were used for behavioral experiments, and males and females were used in equal proportions for all other experiments. For all experiments, animals that carried the *CaMKII $\alpha$ -tTA*, *tetO-PME*, or *tetO-LCMT* transgenes alone were used as controls. We observed no differences among these control groups, suggesting that the phenotypes observed in the *CaMKII $\alpha$ -tTA/tetO-PME* or *CaMKII $\alpha$ -tTA/tetO-LCMT* double-transgenic animals were dependent on *PME-1* or *LCMT-1* transgene expression and were not an artifact of transgene insertion.

**Oligonucleotide in Situ Hybridization and Immunohistochemistry.** RNA in situ hybridization was carried out as described previously (47) using a probe specific to the 3' UTR of both the *LCMT-1* and *PME-1* transgenes. Immunohistochemistry was carried out on sections from paraformaldehyde-perfused animals using an anti-FLAG primary antibody and imaged by confocal microscopy.

**Western Blots.** Western blots were carried out on hippocampal homogenates prepared from microwave-fixed or snap-frozen samples and probed with the indicated antibodies as described previously (13, 48) and as described in more detail in *SI Methods*.

**A $\beta$  Preparation and Infusion.** Oligomeric A $\beta$  was prepared from synthetic A $\beta$  1–42 peptides (American Peptide) as described previously (33, 49). For behavioral experiments, A $\beta$  was infused into the hippocampus via bilaterally implanted cannulae at the indicated volumes and concentrations.

**Fear-Conditioning and Water Maze Tasks.** Contextual fear-conditioning and radial arm water maze tasks were carried out as described previously (27, 30). In the contextual fear-conditioning task freezing behavior was assessed during re-exposure to the training context 24 h after a single foot shock. In the radial arm water maze task, entries into non-platform-containing arms were scored as errors during 10 blocks of three trials, each conducted over 2 d. A single A $\beta$  or vehicle infusion was administered 20 min before training in the contextual fear-conditioning task and 20 min before and midway through training on both days of the radial arm water maze task.

**Electrophysiological Recordings.** Field excitatory postsynaptic potential (fEPSP) recordings of synaptic responses at Schaffer collateral synapses were carried out in 400- $\mu$ m acute hippocampal slices maintained in an interface chamber at 29 °C as described previously (33).

**A $\beta$  ELISA Measures.** A $\beta$ <sub>1–40</sub> and A $\beta$ <sub>1–42</sub> levels were determined by commercially available ELISA kits (Wako) in hippocampal homogenates prepared as described previously (50).

**ACKNOWLEDGMENTS.** We thank Peter Davies (Albert Einstein College of Medicine) for the gift of the PHF-1 antibody. This work was supported by National Institutes of Health Grant 1R01NS092045-01 and Department of Defense Grant W81XWH-12-1-0579 (to O.A. and R.E.N.), a generous gift from the Broitman family (to E.R.K.), and National Health and Medical Research Council of Australia Grant APP1002744 and Hunter Medical Research Institute Grants G1401436 and G1501390 (to E.S. and J.-M.S.).

- Martin L, et al. (2013) Tau protein phosphatases in Alzheimer's disease: The leading role of PP2A. *Ageing Res Rev* 12(1):39–49.
- Sontag JM, Sontag E (2014) Protein phosphatase 2A dysfunction in Alzheimer's disease. *Front Mol Neurosci* 7(16):1–16.
- Voronkov M, Braithwaite SP, Stock JB (2011) Phosphoprotein phosphatase 2A: A novel druggable target for Alzheimer's disease. *Future Med Chem* 3(7):821–833.
- Dreves G, et al. (1993) Dephosphorylation of tau protein and Alzheimer paired helical filaments by calcineurin and phosphatase-2A. *FEBS Lett* 336(3):425–432.
- Gong CX, Grundke-Iqbal I, Iqbal K (1994) Dephosphorylation of Alzheimer's disease abnormally phosphorylated tau by protein phosphatase-2A. *Neuroscience* 61(4):765–772.
- Sontag E, Nunbhakdi-Craig V, Lee G, Bloom GS, Mumby MC (1996) Regulation of the phosphorylation state and microtubule-binding activity of Tau by protein phosphatase 2A. *Neuron* 17(6):1201–1207.
- Xu Y, Chen Y, Zhang P, Jeffrey PD, Shi Y (2008) Structure of a protein phosphatase 2A holoenzyme: Insights into B55-mediated Tau dephosphorylation. *Mol Cell* 31(6):873–885.
- Janssens V, Longin S, Goris J (2008) PP2A holoenzyme assembly: In cauda venenum (the sting is in the tail). *Trends Biochem Sci* 33(3):113–121.
- Sents W, Ivanova E, Lambrecht C, Haesen D, Janssens V (2013) The biogenesis of active protein phosphatase 2A holoenzymes: A tightly regulated process creating phosphatase specificity. *FEBS J* 280(2):644–661.
- Smith AD (2008) The worldwide challenge of the dementias: A role for B vitamins and homocysteine? *Food Nutr Bull* 29(2, Suppl):S143–S172.
- Zhuo JM, Wang H, Praticò D (2011) Is hyperhomocysteinemia an Alzheimer's disease (AD) risk factor, an AD marker, or neither? *Trends Pharmacol Sci* 32(9):562–571.
- Sontag E, et al. (2004) Downregulation of protein phosphatase 2A carboxyl methylation and methyltransferase may contribute to Alzheimer disease pathogenesis. *J Neuropathol Exp Neurol* 63(10):1080–1091.
- Sontag E, et al. (2007) Protein phosphatase 2A methyltransferase links homocysteine metabolism with tau and amyloid precursor protein regulation. *J Neurosci* 27(11):2751–2759.
- Sontag JM, et al. (2008) Folate deficiency induces in vitro and mouse brain region-specific downregulation of leucine carboxyl methyltransferase-1 and protein phosphatase 2A B(alpha) subunit expression that correlate with enhanced tau phosphorylation. *J Neurosci* 28(45):11477–11487.
- Bryant JC, Westphal RS, Wadzinski BE (1999) Methylated C-terminal leucine residue of PP2A catalytic subunit is important for binding of regulatory Balpha subunit. *Biochem J* 339(Pt 2):241–246.
- Longin S, et al. (2007) Selection of protein phosphatase 2A regulatory subunits is mediated by the C terminus of the catalytic subunit. *J Biol Chem* 282(37):26971–26980.
- Ogris E, Gibson DM, Pallas DC (1997) Protein phosphatase 2A subunit assembly: The catalytic subunit carboxy terminus is important for binding cellular B subunit but not polyomavirus middle tumor antigen. *Oncogene* 15(8):911–917.
- Tolstykh T, Lee J, Vafai S, Stock JB (2000) Carboxyl methylation regulates phosphoprotein phosphatase 2A by controlling the association of regulatory B subunits. *EMBO J* 19(21):5682–5691.
- Wei H, et al. (2001) Carboxymethylation of the PP2A catalytic subunit in *Saccharomyces cerevisiae* is required for efficient interaction with the B-type subunits Cdc55p and Rts1p. *J Biol Chem* 276(2):1570–1577.
- De Baere I, et al. (1999) Purification of porcine brain protein phosphatase 2A leucine carboxyl methyltransferase and cloning of the human homologue. *Biochemistry* 38(50):16539–16547.
- Lee J, Stock J (1993) Protein phosphatase 2A catalytic subunit is methyl-esterified at its carboxyl terminus by a novel methyltransferase. *J Biol Chem* 268(26):19192–19195.
- Stanevich V, et al. (2011) The structural basis for tight control of PP2A methylation and function by LCMT-1. *Mol Cell* 41(3):331–342.
- Lee J, Chen Y, Tolstykh T, Stock J (1996) A specific protein carboxyl methyltransferase that demethylates phosphoprotein phosphatase 2A in bovine brain. *Proc Natl Acad Sci USA* 93(12):6043–6047.
- Ogris E, et al. (1999) A protein phosphatase methyltransferase (PME-1) is one of several novel proteins stably associating with two inactive mutants of protein phosphatase 2A. *J Biol Chem* 274(20):14382–14391.
- Xing Y, et al. (2008) Structural mechanism of demethylation and inactivation of protein phosphatase 2A. *Cell* 133(1):154–163.
- Mayford M, et al. (1996) Control of memory formation through regulated expression of a CaMKII transgene. *Science* 274(5293):1678–1683.
- Fiorito J, et al. (2013) Synthesis of quinoline derivatives: Discovery of a potent and selective phosphodiesterase 5 inhibitor for the treatment of Alzheimer's disease. *Eur J Med Chem* 60:285–294.
- Maren S, Phan KL, Liberzon I (2013) The contextual brain: Implications for fear conditioning, extinction and psychopathology. *Nat Rev Neurosci* 14(6):417–428.
- Watterson DM, et al. (2013) Development of novel in vivo chemical probes to address CNS protein kinase involvement in synaptic dysfunction. *PLoS One* 8(6):e66226.
- Alamed J, Wilcock DM, Diamond DM, Gordon MN, Morgan D (2006) Two-day radial-arm water maze learning and memory task; robust resolution of amyloid-related memory deficits in transgenic mice. *Nat Protoc* 1(4):1671–1679.
- Ondrejcek T, et al. (2010) Alzheimer's disease amyloid beta-protein and synaptic function. *Neuromolecular Med* 12(1):13–26.
- Haass C, Selkoe DJ (2007) Soluble protein oligomers in neurodegeneration: Lessons from the Alzheimer's amyloid beta-peptide. *Nat Rev Mol Cell Biol* 8(2):101–112.
- Puzzo D, et al. (2008) Picomolar amyloid-beta positively modulates synaptic plasticity and memory in hippocampus. *J Neurosci* 28(53):14537–14545.
- Lambert MP, et al. (1998) Diffusible, nonfibrillar ligands derived from Abeta1–42 are potent central nervous system neurotoxins. *Proc Natl Acad Sci USA* 95(11):6448–6453.
- Puzzo D, Arancio O (2013) Amyloid- $\beta$  peptide: Dr. Jekyll or Mr. Hyde? *J Alzheimers Dis* 33(Suppl 1):S111–S120.
- Puzzo D, et al. (2011) Endogenous amyloid- $\beta$  is necessary for hippocampal synaptic plasticity and memory. *Ann Neurol* 69(5):819–830.
- Yu XX, et al. (2001) Methylation of the protein phosphatase 2A catalytic subunit is essential for association of Balpha regulatory subunit but not SG2NA, striatin, or polyomavirus middle tumor antigen. *Mol Biol Cell* 12(1):185–199.
- Matsuo ES, et al. (1994) Biopsy-derived adult human brain tau is phosphorylated at many of the same sites as Alzheimer's disease paired helical filament tau. *Neuron* 13(4):989–1002.
- Longin S, et al. (2004) An inactive protein phosphatase 2A population is associated with methyltransferase and can be re-activated by the phosphotyrosyl phosphatase activator. *Biochem J* 380(Pt 1):111–119.
- Longin S, et al. (2008) Spatial control of protein phosphatase 2A (de)methylation. *Exp Cell Res* 314(1):68–81.
- Wandzioch E, et al. (2014) PME-1 modulates protein phosphatase 2A activity to promote the malignant phenotype of endometrial cancer cells. *Cancer Res* 74(16):4295–4305.
- Dawkins E, Small DH (2014) Insights into the physiological function of the  $\beta$ -amyloid precursor protein: Beyond Alzheimer's disease. *J Neurochem* 129(5):756–769.
- Müller UC, Zheng H (2012) Physiological functions of APP family proteins. *Cold Spring Harb Perspect Med* 2(2):a006288.
- Parihar MS, Brewer GJ (2010) Amyloid- $\beta$  as a modulator of synaptic plasticity. *J Alzheimers Dis* 22(3):741–763.
- Lambrecht C, Haesen D, Sents W, Ivanova E, Janssens V (2013) Structure, regulation, and pharmacological modulation of PP2A phosphatases. *Methods Mol Biol* 1053:283–305.
- Committee on Care and Use of Laboratory Animals (1996) *Guide for the Care and Use of Laboratory Animals* (Natl Inst Health, Bethesda), DHHS Publ No (NIH) 85-23.
- Wisden W, Morris BJ (2002) In situ hybridization with oligonucleotide probes. *Int Rev Neurobiol* 47:3–59.
- Bottiglieri T, et al. (2012) Acute administration of L-DOPA induces changes in methylation metabolites, reduced protein phosphatase 2A methylation, and hyperphosphorylation of Tau protein in mouse brain. *J Neurosci* 32(27):9173–9181.
- Stine WB, Jr, Dahlgren KN, Krafft GA, LaDu MJ (2003) In vitro characterization of conditions for amyloid-beta peptide oligomerization and fibrillogenesis. *J Biol Chem* 278(13):11612–11622.
- Teich AF, Patel M, Arancio O (2013) A reliable way to detect endogenous murine  $\beta$ -amyloid. *PLoS One* 8(2):e55647.
- Lois C, Hong EJ, Pease S, Brown EJ, Baltimore D (2002) Germline transmission and tissue-specific expression of transgenes delivered by lentiviral vectors. *Science* 295(5556):868–872.
- Lee L, Kosuri P, Arancio O (2014) Picomolar amyloid- $\beta$  peptides enhance spontaneous astrocyte calcium transients. *J Alzheimers Dis* 38(1):49–62.
- Paxinos G, Franklin KJB (2004) *The Mouse Brain in Stereotaxic Coordinates* (Elsevier Academic, Amsterdam) Compact 2nd Ed.
- Arning E, Bottiglieri T (2016) Quantitation of S-adenosylmethionine and S-adenosylhomocysteine in plasma using liquid chromatography-electrospray tandem mass spectrometry. *Methods Mol Biol* 1378:255–262.